arXiv:1004.0266v1 [hep-th] 2 Apr 2010

A dynamical symmetry for supermembranes

JONAS DE WOUL[1,*], JENS HOPPE[2,†], DOUGLAS LUNDHOLM[2,‡], AND MARTIN SUNDIN[1,§]

[1] *Department of Theoretical Physics, Royal Institute of Technology (KTH)*
106 91 Stockholm, Sweden

[2] *Department of Mathematics, Royal Institute of Technology (KTH)*
100 44 Stockholm, Sweden

Abstract

A dynamical symmetry for supersymmetric extended objects is given.

Applying the Noether procedure to the supermembrane action [1] in a light-cone formulation, and using work of Goldstone [2], who proved Lorentz invariance of the corresponding bosonic theory by finding a way to explicitly reconstruct ζ (usually denoted x^-) in terms of the transverse degrees of freedom, the generators of "mixed" rotations (involving the corresponding supersymmetrised version of ζ) were given in [3] (as expressions involving $\vec{x}$, $\vec{p}$, θ, and, as in the bosonic theory, two discrete degrees of freedom: η and $\zeta_0 = \int \rho\zeta d^2\varphi$) as

$$J_{i-} = \int \Big(x_i \mathcal{H}_S - \zeta_S p_i - \frac{\mathrm{i}}{4\eta}\theta\gamma^{ik}\theta p_k - \frac{\mathrm{i}}{8\eta}\epsilon^{rs}\partial_r x_j \partial_s x_k \theta\gamma^{ijk}\theta \Big) d^2\varphi, \tag{1}$$

with

$$\zeta_S = \zeta_0 - \frac{1}{\eta}\int G^r(\varphi,\tilde{\varphi})\Big(\frac{\vec{p}}{\rho}\cdot\tilde{\partial}_r\vec{x} + \frac{i}{2}\theta\tilde{\partial}_r\theta\Big)\rho(\tilde{\varphi})d^2\tilde{\varphi}, \tag{2}$$

$$\mathcal{H}_S = \frac{\vec{p}^2 + \det(\partial_r\vec{x}\cdot\partial_s\vec{x})}{2\eta\rho} - \frac{\mathrm{i}}{2\eta}\theta\gamma^i\epsilon^{rs}\partial_r x_i \partial_s\theta, \tag{3}$$

$$\int G^r(\varphi,\tilde{\varphi})\rho d^2\varphi = 0, \quad \tilde{\nabla}_r G^r(\varphi,\tilde{\varphi}) = \frac{\delta(\varphi,\tilde{\varphi})}{\rho} - 1, \tag{4}$$

the $\vec{x}_i$, $\vec{p_j}$ canonically conjugate, and

$$\{\theta_\alpha(\varphi), \theta_\beta(\tilde{\varphi})\} = -\frac{\mathrm{i}}{\rho}\delta_{\alpha\beta}\delta(\varphi,\tilde{\varphi}), \quad \{\eta, \zeta_0\} = 1, \tag{5}$$

(all other dynamical Poisson brackets zero).

---

[*] jodw02@kth.se
[†] hoppe@math.kth.se
[‡] dogge@math.kth.se
[§] masundi@kth.se

Recently, one of us [4] found a dynamical symmetry for (bosonic) relativistic extended objects of any dimensionality. In the present note, it is pointed out that an analogous dynamical symmetry exists for supermembranes. The generators of internal transverse rotations,

$$\bar{J}_{ij} = \int \Big(x_i p_j - x_j p_i - \frac{\mathrm{i}}{4}\theta\gamma^{ij}\theta\rho \Big) d^2\varphi - \Big(X_i P_j - X_j P_i \Big) + \frac{\mathrm{i}}{4}\theta_0\gamma^{ij}\theta_0, \tag{6}$$

$$X_i := \int \rho x_i d^2\varphi, \quad P_i := \int p_i d^2\varphi, \quad \theta_0 := \int \rho\theta d^2\varphi, \tag{7}$$

and the purely internal (non zero mode) part of (1), denoted by $\bar{J}_{i-}$, together with an additional (new) angular momentum formed out of dynamically generated Clifford variables, constitute a super-generalisation of the dynamical symmetry obtained in [4]. Assuming that

$$\{J_{i-}, J_{k-}\} = 0, \tag{8}$$

the dynamical Poisson brackets $\{\bar{J}_{i-}, \bar{J}_{k-}\}$ can be inferred by subtracting from (1) all terms involving zero modes:

$$\bar{J}_{i-} := J_{i-} - J^{(0)}_{i-} - \tilde{J}_{i-}, \tag{9}$$

$$J^{(0)}_{i-} := \Big(X_i H_S - \zeta_0 P_i \Big) - \frac{\mathrm{i}}{4\eta}\theta_0\gamma^{ik}\theta_0 P_k, \tag{10}$$

$$\tilde{J}_{i-} := \frac{1}{\eta}\bar{J}_{ik}P_k - \frac{\mathrm{i}}{2\eta}\theta_0\gamma^i Q, \tag{11}$$

$$Q_\beta := \int \Big(p_i \gamma^i_{\beta\alpha}\theta_\alpha + \frac{1}{2}\epsilon^{rs}\partial_r x_i \partial_s x_j \gamma^{ij}_{\beta\alpha}\theta_\alpha \Big) d^2\varphi - P_i\gamma^i_{\beta\alpha}\theta_{0\alpha}. \tag{12}$$

On the physical phase space, constrained by

$$\epsilon^{rs}\Big(\partial_r \vec{p} \cdot \partial_s \vec{x} + \frac{\mathrm{i}}{2}\partial_r\theta\partial_s\theta \Big) = 0, \tag{13}$$

we find

$$\{J^{(0)}_{i-} + \tilde{J}_{i-}, J^{(0)}_{j-} + \tilde{J}_{j-}\} = -\frac{\mathbb{M}^2}{\eta^2}\bar{J}_{ij} - \frac{\mathrm{i}}{4\eta^2}Q_\alpha\gamma^{ij}_{\alpha\beta}Q_\beta, \tag{14}$$

with $\mathbb{M}^2$ the relativistically-invariant internal (squared) mass

$$\mathbb{M}^2 = 2\eta \int \mathcal{H}_{\mathcal{S}} d^2\varphi - \vec{P}^2. \tag{15}$$

Assuming that

$$\{\bar{J}_{i-}, Q_\beta\} = 0 \tag{16}$$

(which we deduced from a general argument, but better be checked explicitly as well), the cross terms vanish

$$\{\bar{J}_{i-}, J^{(0)}_{j-} + \tilde{J}_{j-}\} + \{J^{(0)}_{i-} + \tilde{J}_{i-}, \bar{J}_{j-}\} = 0. \tag{17}$$

Hence we deduce that

$$\{\eta\bar{J}_{i-}, \eta\bar{J}_{k-}\} = \mathbb{M}^2\bar{J}_{ik} - Q_{ik}, \tag{18}$$

with

$$Q_{ik} := -\frac{\mathrm{i}}{4} Q_\alpha \gamma^{ik}_{\alpha\beta} Q_\beta. \tag{19}$$

For $\mathbb{M}^2 \neq 0$, one could also write this as

$$\{\mathbb{M}_i, \mathbb{M}_k\} = \bar{J}_{ik} - \Sigma_{ik}, \tag{20}$$

where

$$\mathbb{M}_i := \frac{\eta \bar{J}_{i-}}{\sqrt{\mathbb{M}^2}}, \tag{21}$$

and

$$\Sigma_{ik} := -\frac{\mathrm{i}}{4} \frac{Q_\alpha}{\sqrt{\mathbb{M}^2}} \gamma^{ik}_{\alpha\beta} \frac{Q_\beta}{\sqrt{\mathbb{M}^2}}, \tag{22}$$

is formed out of dynamically generated Clifford variables

$$\Sigma_\beta := \frac{Q_\beta}{\sqrt{\mathbb{M}^2}}, \quad \{\Sigma_\alpha, \Sigma_\beta\} = \delta_{\alpha\beta}. \tag{23}$$

Observing that

$$\begin{aligned}
\{Q_{ij}, Q_{kl}\} &= \Big(-\delta_{jk} Q_{il} + \delta_{ik} Q_{jl} - \delta_{il} Q_{jk} + \delta_{jl} Q_{ik}\Big)\mathbb{M}^2, && (24)\\
\{\bar{J}_{ik}, \eta \bar{J}_{l-}\} &= -\delta_{kl} \eta \bar{J}_{i-} + \delta_{il} \eta \bar{J}_{k-}, && (25)\\
\{Q_{ik}, \eta \bar{J}_{l-}\} &= 0, && (26)\\
\{\bar{J}_{ij}, Q_{kl}\} &= -\delta_{jk} Q_{il} + \delta_{ik} Q_{jl} - \delta_{il} Q_{jk} + \delta_{jl} Q_{ik}, && (27)
\end{aligned}$$

one finally notes that, assuming (8) and (16), $\eta \bar{J}_{i-}$ and $M_{ik} := \mathbb{M}^2 \bar{J}_{ik} - Q_{ik}$ (commuting with the supercharges Q_α) generate the following dynamical symmetry algebra for supermembranes

$$\begin{aligned}
\{M_{ij}, M_{kl}\} &= \Big(-\delta_{jk} M_{il} + \delta_{ik} M_{jl} - \delta_{il} M_{jk} + \delta_{jl} M_{ik}\Big)\mathbb{M}^2, && (28)\\
\{\eta \bar{J}_{i-}, \eta \bar{J}_{k-}\} &= M_{ik}, && (29)\\
\{M_{ik}, \eta \bar{J}_{l-}\} &= \Big(-\delta_{kl} \eta \bar{J}_{i-} + \delta_{il} \eta \bar{J}_{k-}\Big)\mathbb{M}^2. && (30)
\end{aligned}$$

Acknowledgement

We would like to thank E. Langmann for his kind support.

References

[1] E. Bergshoeff, E. Sezgin, P. K. Townsend, Phys. Lett. B. **189**, 75 (1987)

[2] J. Goldstone, unpublished notes

[3] B. de Wit, U. Marquard, H. Nicolai, Commun. Math. Phys. **128**, 39 (1990)

[4] J. Hoppe, `arXiv:1003.5189`